



Understanding Cataracts



For more information, please contact

T 0116 249 8839

E info@vistablind.org.uk

W www.vistablind.org.uk

Contents


- 3** About cataracts
- 4** How your eye works
- 5** Causes
- 7** Symptoms
- 9** Treatment
- 19** Complications
- 21** Other eye conditions
- 21** Coping
- 23** Useful contacts
- 23** About Vista

About cataracts

Cataracts are a very common eye condition. As we get older, the lens inside our eye gradually changes and becomes less transparent (clear). A lens that has turned misty or cloudy is said to have a cataract. Over time a cataract can get worse, gradually making your vision mistier.

A straightforward operation can usually remove the misty lens and replace it with an artificial lens to enable you to see more clearly again.

This leaflet is about cataracts in adults. Some children develop cataracts, called congenital cataracts, before or just after birth but these are usually dealt with in a different way to cataracts in adults.



How your eye works

When you look at something, light passes through the front of the eye and is focused by the cornea and then the lens onto the retina. The lens is normally clear so that light can pass directly through to focus on the retina (the lens is clear because of the way the cells in the lens are arranged).

The lens focusses light onto the retina, which converts the light into electrical signals. A network of nerves delivers these signals from the different parts of the retina to the optic nerve and then onto the brain. The brain interprets these signals to 'see' the world around us.

The lens can change shape, allowing us to focus on objects at different distances, called 'accommodation of vision'. As we get older, the lens isn't able to change shape as well as it used to; even people who can see clearly in the distance without glasses will need reading glasses to see things up close. This process is not caused by a cataract.

Cataracts result from changes in the way cells of the lens are arranged and their water content, which causes the lens to become cloudy instead of clear. When this happens, light cannot pass directly through the lens and you may notice problems with your vision. A cataract is not a growth or a film growing over the eye; it is simply the lens becoming misty.

Causes

Cataracts can be caused by a number of things, but by far the most common reason is growing older. Most people over the age of 65 have some changes in their lens and most of us will develop a cataract in time. Apart from getting older, the other common causes of cataracts include:

- Diabetes
- Trauma
- Medications, such as steroids
- Eye surgery for other eye conditions
- Other eye conditions.

In general, the reason why you have developed a cataract will not affect the way it is removed.

Most cataracts are caused by natural changes in the lens, which happens as you get older. However, the following factors may be involved in cataract development (please note that these are only suggested causes which are the subject of ongoing research):

- Tobacco smoking
- Lifelong exposure to sunlight
- Having a poor diet lacking antioxidant vitamins.

Symptoms

Cataracts usually develop slowly, and although symptoms vary there are some symptoms that most people experience. Most people will eventually develop a cataract in both eyes, though one eye may be affected before the other. When your cataract starts to develop, you may feel your sight isn't quite right. For example, if you wear glasses you may feel that your lenses are dirty, even when they're clean.

Gradually, you may find your sight becomes cloudier and more washed out. Edges of stairs or steps become more difficult to see and you may feel you need a bit more light to read smaller print.

Another common symptom of a cataract is a problem with bright lights. Lights can seem to glare or you may find that the headlights of a car dazzle you more than they used to. You may also notice a slight change in your colour vision – things may appear more yellow than before. This often happens if one eye develops a cataract first and colours look different when you compare one eye with the other.

If you notice any of these changes, you should have your eyes tested by an optometrist (optician) who will be able to tell whether you have a cataract or not.

The optometrist will then discuss the degree to which the cataract is affecting your vision and, if you agree, refer you via your GP to the eye clinic.

You may be told during the eye test that you have an early cataract or initial signs of a cataract which does not need referral.

If you are unsure about anything during the eye test then ask to have it explained.

Eye tests are free for everyone in Scotland and for everyone over 60 years old in the UK. Many other people also qualify for free eye tests. If you are unsure speak to your optometrist.

If a cataract isn't removed, your sight will become increasingly cloudy. Eventually, it will be like trying to see through a frosted window or a heavy set curtain of fog. Even if your cataract gets to this stage, it can be removed and your sight will be almost as it was before the cataract developed.

Treatment

The only affective treatment for cataracts is surgery to remove your cloudy lens and replace it with an artificial lens implant. This is done by an ophthalmologist (eye specialist) at a hospital.

Lasers are not used to remove cataracts, and there is no evidence to suggest that changing your diet, taking vitamins or using eye drops can cure cataracts. Cataract surgery is available free on the NHS.

Removing cataracts

The operation to remove your cataracts can be performed at any stage of their development.

There is no longer a reason to wait until your cataract is 'ripe' before removing it. However, because any surgery involves some risk, it is usually worth waiting until there is some change in your vision before removing the cataract. This is something to discuss with your optometrist as a good time to refer may vary from person to person.

Most people choose to have their cataracts removed when their change in vision starts to cause them difficulties in everyday life. The timing of this varies from person to person. If you have problems in bright light, or you find reading or getting out and about, cooking or looking after yourself increasingly difficult, then it may be time to consider having the cataract removed.

When you attend your appointment in the eye clinic, you need to make clear to the specialists any everyday problems you are having.

When you are first referred to the eye clinic, you will have an outpatient appointment to examine your eyes and then discuss the best options for you. This is the time to ask questions and it is useful to write down any you have thought of beforehand.

Many people with cataracts are still legally able to drive. If you have any concerns about whether you should be driving, your optometrist should be able to tell you whether your sight is within the legal limits for driving. Sometimes people may be legally able to drive but might find driving difficult in bright sunlight or at night. If this is the case, then you may think it is a good time to consider having your cataracts removed.

Pre-surgery

Before you have your cataract surgery, your eye health and general health will be checked carefully in what is often called a pre-operative assessment. Your vision and your eye will be measured very carefully.

This is usually done by a machine, which measures the length of your eye ball and the shape of the front of your eye. These tests help the ophthalmologist to decide which lens to implant when they perform your operation, and to make sure your vision is as good as possible after the surgery.

If you have a cataract in both eyes, your ophthalmologist will use these tests to decide which cataract to remove in the first operation. In most cases, this is the eye with the worst cataract.

Surgery

Cataract surgery usually takes about 30 to 40 minutes, and most people go home from hospital a few hours later. It is usually done with a local anaesthetic, which means you will be awake during the operation but you won't feel any pain. You can talk to the operating team if you need any assurance. The local anaesthetic may involve drops and/or an injection.

For your surgery, you will be given drops to dilate your pupil. Your face will be covered by a sheet, which helps to keep the area around your eye clean during the operation.

To remove the cataract, the ophthalmologist needs to remove the natural lens in your eye and replace it with a plastic lens implant. The most common way to remove cataracts is called Phacoemulsification.

This technique uses high frequency sound energy to break up your natural lens with the cataract. Only really small cuts are used, so you don't need any stitches, and this helps to speed up your recovery from the surgery. Usually, the ophthalmologist uses a machine which acts as a microscope to get the best view of your eye as possible.

The lens in your eye is made up of different layers and the outside layer is called the lens capsule. During the operation, the ophthalmologist cuts through the front of the lens capsule so they can reach the lens inside. Using the same instrument, the ophthalmologist can break up your lens and the cataract inside your eye and remove it using suction.

Your lens capsule is kept in place so that the artificial lens implant can be placed inside it. The tiny implant is folded, so that it can be put into the eye through the same instrument that is used to remove the cataract. Once it reaches the right position, the ophthalmologist unfolds the artificial lens so that it sits in the right place inside the lens capsule.

As you are awake during the operation, you will be able to hear what is happening in the operating room. You can also communicate with the ophthalmologist and the nurses who are on hand to reassure you.

Because the eye is anaesthetised and the pupil is dilated, you may be able to see some lights and movement but not the details of the instruments used. You should not feel any pain in your eye.

A short time after your operation, your eye will be examined to make sure the operation has been a success. Your eye will be covered with a dressing which stays in place when you go home, normally a few hours later.

Your eye may begin to feel sore once the anaesthetic wears off. The pain isn't usually too bad and you can take a painkiller tablet, such as paracetamol, to help. The dressing, which is put on in the hospital, usually needs to stay on your eye overnight, but you should be able to take it off the following morning.

Your eye may look red and you might develop some bruises, but these will improve over the next few days.

Immediately after the operation

Some people can tell that their sight has improved straight away. If your cataract was quite mild, you might not notice a big change in your vision, but if your cataract was quite bad, you may be able to notice a bigger improvement. Either way, your sight may not be as good as you expected for the first week after the operation, as the eye is still recovering from the surgery and will probably be a little swollen.

Immediately after the surgery you will be given eye drops. You will probably have two types of drops – an antibiotic drop to prevent infection and a steroid drop to help reduce the swelling. It is important to take these drops as the ophthalmologist recommends and to complete the course. If you have problems using the drops you should let your GP know as they may be able to arrange some help for you.

Most people have no problems after the surgery and they are up and about as normal the next day. If your eye is very painful or your vision suddenly gets a lot worse, then you should let the hospital know as soon as possible, as this may mean they need to see you again.

Activities

After surgery, you can usually go back to your everyday activities as soon as you feel able. Apart from taking eye drops, you can usually carry on as normal, but you may need to avoid the following for the first week to ten days:

- Rubbing your eye. You may have to wear an eye shield (patch) when you are sleeping to avoid rubbing your eye
- Swimming (until your ophthalmologist says you can). You should avoid contact with dirty water whilst your eye is healing
- Strenuous exercise, contact sports and heavy lifting. Everyday lifting like light shopping is usually fine, but heavy lifting like moving furniture is best avoided
- Wearing eye make up until the hospital is happy with your recovery.

You also need to take care:

- When it is windy or dusty outdoors, in case something blows in your eye, but you don't need to stay indoors
- Washing your hair - avoid soapy water in your eye.

When should I have new glasses?

The lens that is implanted in your eye is usually designed to give you clear distance vision without needing glasses. Sometimes this is not quite achieved and you will need a pair of distance glasses to fine-tune the focus and get the best possible distance vision. Because the lens implant isn't able to provide in-focus near vision, nearly everyone needs to wear reading glasses after the operation and usually this is a different pair than you had before the operation.

In most cases an eye test, sometimes called a refraction, will be done four to six weeks after the operation. This may be done by an optometrist in the hospital or you may be asked to see your own optometrist.

Some lens implants are available which try to provide clear vision in the distance and close up. These are called multi-focal lenses. There are different types available, but they are usually implanted in the same way as the more common lenses. At the moment, multi-focal lenses are normally only available privately.

Between operations

If you have cataracts in both eyes, the period between having the first and second operation can be difficult. This is because your eye will not be balanced in terms of glasses and correction for any short or long sight you may have.

Normally, people are encouraged to wait until they have a second operation before getting new glasses. This avoids the need to buy glasses that would only be useful for the short period between the operations. Some people find they can manage with their old reading glasses, but this may not be possible for everyone. The gap between the two operations is usually six weeks to three months so most people can manage.

Complications

Cataract surgery is generally very successful. Only about three percent of people who have cataracts experience complications. The most common complications can be dealt with and usually don't affect sight in the long term.

One of the most common complications is thickening of the lens capsule, which holds the lens in place. This may occur a couple of months or even years after the original operation.

If this happens, your sight will become cloudy again, as though the cataract has come back. Doctors call this complication posterior capsule opacification or posterior capsule thickening, and it is usually dealt with by a small laser operation done through an outpatient appointment.

More serious complications are much rarer and include:

- Retinal detachment
- Problems with the lens implant, the wiring lens implant or problems with its position
- A break in the lens capsule
- Infection.

These complications are much rarer and treatments are available, which will minimise their effects on your vision. Before being offered a cataract operation, the ophthalmologist will talk you through the potential risks specific to your situation.

Other eye conditions

People with cataracts often have other eye conditions as well. This is because many eye conditions affect older people. For example, many people with macular degeneration, glaucoma or diabetic eye problems also develop cataracts.

Removing cataracts when you have other eye conditions is possible, but there may be other things to consider. Ask your ophthalmologist to explain any extra considerations for a cataract operation in your particular circumstances.

Coping

Being diagnosed with an eye condition can be very upsetting. You may find that you are worried about the future and how you will manage with a change in your vision. All these feelings are natural.

Some people may want to talk over some of these feelings with someone outside their circle of friends or family. Vista can help, with our telephone helpline and our emotional support service. Your GP or social worker may also be able to help you find a counsellor if you think this would help you.

Most people who have cataracts without any other eye condition have very good vision following the operation. However, if you have another eye condition that affects your sight, or there are complications with the cataract surgery which affect your sight, then there are lots of things you can do to make the most of the vision you have.

This may mean making things bigger, brighter or using colour to help you see things better. Ask your ophthalmologist, optician or GP to refer you to our Low Vision Clinic.

The Vista helpline can also give you information about the low vision service and the local support that is available either through Vista or social services.

Call us on **0116 249 8839** or visit our website at **www.vistablind.org.uk**.

Useful contacts

The Royal College of Ophthalmologists

T 020 7935 0702

W www.rcophth.ac.uk

Royal National Institute of Blind People (RNIB)

T 0303 123 9999

W www.rnib.org.uk

Driver and Vehicle Licensing Agency (DVLA)

T 0300 790 6801

W www.gov.uk/contact-the-dvla

About Vista

Vista is the leading local charity working with children and adults with sight loss in Leicester, Leicestershire and Rutland. Whether someone is newly diagnosed or has lived their whole life with reduced sight, Vista's specialist staff provide a vital lifeline to local people.

T 0116 249 8839

E info@vistablind.org.uk

W www.vistablind.org.uk



The information in this leaflet has been reproduced from the booklet “Understanding Cataracts” with kind permission of the Royal College of Ophthalmologists and RNIB.

If you require the information in this booklet in an alternative format (e.g. large print, audio or Braille) or in an alternative language, please contact Vista on **0116 249 0909.**

For more information, please contact

Vista

16 New Walk
Leicester
LE1 6UB

T 0116 249 8839

E info@vistablind.org.uk

www.vistablind.org.uk

Registered charity number 218992